

Observational Tests of Instanton Cosmology

R.E. Allen, J. Dumoit, and A. Mondragon

Center for Theoretical Physics, Texas A&M University

College Station, Texas, 77843 USA

e-mail: allen@tamu.edu

A new cosmological model leads to testable predictions that are different from those of both standard cosmology and models with a cosmological constant. The prediction that $q_0 = 0$ is the same as in other “coasting universe” models, but arises without the need for any exotic form of matter or other *ad hoc* assumptions.

A new cosmological model was recently proposed, in which the big-bang singularity and global metric of spacetime are due to a cosmological instanton.^{1,2,3} The starting point is the order parameter Ψ_s for a GUT Higgs field which has a symmetry group $\text{SO}(10) \times \text{SU}(2) \times \text{U}(1)$.¹ Since the coupling to other fields is not discussed in the present paper, the $\text{SO}(10)$ gauge symmetry will be ignored. Ψ_s can then be treated simply as an $\text{SU}(2) \times \text{U}(1)$ order parameter, and there are two natural point defects:

(1) $\text{SU}(2)$ instantons, which are the four-dimensional generalization (in Euclidean spacetime) of two-dimensional vortices in ordinary superfluids. With the order parameter written in the form $\Psi_s = n_s^{1/2} U \eta_0$ and the scalings $\rho = r/\xi$, $f = (n_s/\bar{n}_s)^{1/2}$, $\xi = (2m\mu)^{-1/2}$, $\bar{n}_s = \mu/b$ in a standard notation,⁴ the condensate density n_s satisfies the generalized Bernoulli equation^{1,2}

$$-\frac{1}{\rho^3} \frac{d}{d\rho} \left(\rho^3 \frac{df}{d\rho} \right) + \frac{3a^2}{\rho^2} f - f + f^3 = 0 \quad (1)$$

in Euclidean spacetime. The asymptotic solutions are

$$f = 1 - 3a^2/2\rho^2 \quad , \quad \rho \rightarrow \infty \quad \text{and} \quad f = C\rho^n \quad , \quad \rho \rightarrow 0 \quad (2)$$

where C is a constant and $n = (1 + 3a^2)^{1/2} - 1$. An $n = 1$ BPST-like instanton then has a strength $a = 1$.

(2) $\text{U}(1)$ monopole-like defects, which act as sources of condensate current, and which have a Euclidean action^{1,2}

$$S = \bar{n}_s \mu \int d^4x f^\dagger \left[-\frac{1}{\rho^3} \frac{d}{d\rho} \left(\rho^3 \frac{df}{d\rho} \right) - \frac{\bar{a}^6}{\rho^6 (f^\dagger f)^2} f - f + \frac{1}{2} (f^\dagger f) f \right]. \quad (3)$$

In the Euclidean formulation, f is not necessarily real and $f^\dagger$ is not even necessarily the Hermitian conjugate of f . (See the comments below (2.10) and

(3.12) of Ref. 1.) Here, however, we will be concerned only with the behavior of Ψ_s in the Lorentzian spacetime of human observers, where f is real and n_s is positive.

A general point defect can have both a and $\bar{a}$ nonzero. The cosmological instanton of Refs. 1 and 2 is such a defect, and the radial coordinate in four-dimensional spacetime is interpreted as the time coordinate: $x^0 = r$. After a Wick rotation to Lorentzian spacetime, the density of the bosonic field Ψ_s is determined by the equation

$$\frac{1}{\rho^3} \frac{d}{d\rho} \left(\rho^3 \frac{df}{d\rho} \right) + \frac{3a^2}{\rho^2} f - \frac{\bar{a}^6}{\rho^6 f^4} f - f + f^3 = 0 \quad (4)$$

and the asymptotic solutions are

$$f = 1 - 3a^2/2\rho^2 \quad , \quad \rho \rightarrow \infty \quad \text{and} \quad f = c\rho^{-1} \quad , \quad \rho \rightarrow 0. \quad (5)$$

Although f diverges as $\rho \rightarrow 0$, there is a cutoff at the Planck length ℓ_P ,^{1,2} which is analogous to the cutoff at the atomic scale in an ordinary superfluid.

Let ρ_c be defined by $f(\rho_c) = 1$ and let $\bar{\eta} = \bar{a}/a$. Realistic values of $\bar{a}$ and a are, of course, cosmologically large. For example, if $\bar{t}$ is the value of the time t (in the Robertson-Walker metric) that corresponds to $\rho = \bar{a}$, and if $\bar{t}$ is ~ 100 seconds, the results of Ref. 2 imply that $\bar{a} \sim 10^{44}$. Nevertheless, it is informative to solve the differential equation (4) numerically for more modest values, and Fig. 1 shows the solution for $\bar{a} = \bar{\eta} = 10$. One expects that $f(\rho)$ is slowly decreasing for $\bar{a} < \rho < \rho_c$, that $f(\rho) \approx 1$ in this same range, and that $\rho_c = \bar{a} (\bar{a}^2/3a^2)^{1/4}$ to a very good approximation. These expectations are confirmed by the results of Fig. 1, and by calculations that we have performed for other values including $\bar{a} = \bar{\eta} = 100$.

In the present model, the universe is born at $\rho \sim 1$ (or $x^0 \sim t_P \sim 10^{-42}$ sec) with a highly confined and coherent bosonic field Ψ_s . Through interactions with other fields that are omitted in the present paper, it releases a small portion of its initially very large energy, producing a hot big bang. At the same time, it relaxes into a fully condensed state during the period $0 < \rho < \rho_c$. It then remains fully condensed, so that $n_s = \bar{n}_s$ for $\rho \geq \rho_c$. This requires that²

$$v_\alpha^0 v_\alpha^0 = v_\alpha^k v_\alpha^k \quad , \quad \rho > \rho_c \quad (6)$$

where

$$v^\mu = v_\alpha^\mu \sigma^\alpha = -im^{-1} U^{-1} \partial^\mu U \quad (7)$$

is the generalization of $m\vec{v} = \vec{\nabla}\theta$ in an ordinary superfluid.

The condition (6) has the form $\omega^2 = p^2$, where $p^k = mv^k$ describes the rotations of the two-component order parameter Ψ_s as a function of position.




Fig. 1. Scaled value of $n_s^{1/2}$ versus scaled value of x^0 , for $a = 1$ and $\bar{a} = 10$.

This condition can be written in a third form: $v_\alpha^\mu v_\mu^\alpha = \eta_{\mu\nu} v_\alpha^\mu v_\alpha^\nu = 0$, where $\eta_{\mu\nu} = \text{diag}(-1, 1, 1, 1)$ is the Minkowski metric tensor. When it is satisfied, the generalized Bernoulli equation

$$-\frac{1}{2m}n_s^{-1/2}\partial^\mu\partial_\mu n_s^{1/2} + \frac{1}{2}mv_\alpha^\mu v_\mu^\alpha + bn_s = \mu \quad (8)$$

holds with $n_s = \bar{n}_s = \mu/b$. I.e., the GUT Higgs field Ψ_s is fully condensed for large times $\rho > \rho_c$.

In the theory of Ref. 1, the geometry of spacetime is determined by the field v_α^μ , which is interpreted as the vierbein: $g^{\mu\nu} = \eta^{\alpha\beta} e_\alpha^\mu e_\beta^\nu$ with $e_\alpha^\mu = v_\alpha^\mu$. For example, massless fermions and vector bosons move just as in standard general relativity, except for a breaking of Lorentz invariance at high energies, and a small additional gravitational interaction in the case of fermions. The local curvature of spacetime results from Planck-scale instantons, just as the rotation of a superfluid results from quantized vortices. The Einstein field equations hold as a very good approximation for local deformations of spacetime geometry, but the global metric of the universe is determined by the cosmological $SU(2) \times U(1)$ instanton described above. One obtains a metric with the usual Robertson-Walker form

$ds^2 = -dt^2 + R(t)^2 [dr^2/(1-r^2) + r^2 d\Omega]$ but with a cosmic scale factor which is quite different from that of standard Friedmann models:²

$$R(t) = \eta t \quad , \quad t \rightarrow 0 \quad (9)$$

$$R(t) \propto \left[\int dt/n_s(t) \right]^{1/2} \quad , \quad t < t_c \quad (10)$$

$$dR(t)/dt = 2\sqrt{3} \quad , \quad t > t_c \quad (11)$$

where $\eta \approx 2\bar{\eta} \gg 1$. Recall that $n_s(t)$ is a decreasing function of t and $n_s(t) \approx \bar{n}_s$ for $\bar{t} < t < t_c$. These forms follow from the solutions to (4) and (6), with the details given in Ref. 2. Here $\bar{t}$ and t_c are the times corresponding to $\rho = \bar{\rho}$ and $\rho = \rho_c$ respectively. For example, $\bar{t} \sim 100$ sec and $\bar{\eta} \sim 100$ corresponds to $t_c \sim 10^6$ sec.

The behavior of the scale factor in (9)-(11) has major implications:² (9) and (11) offer solutions to the smoothness, monopole, and flatness problems, and (10) is fully consistent with big-bang nucleosynthesis. In addition, (11) implies a comfortably large age for the universe, and predicts that the deceleration parameter q_0 is exactly zero. This is quite different from the predictions $q_0 \geq 1/2$, for Friedmann models with a flat or closed universe, or $q_0 \sim -1/2$, for Λ CDM models.⁵ There have been other proposals for a “coasting universe”, but they involve exotic forms of matter⁶ or other *ad hoc* assumptions,⁷ whereas here the result $q_0 = 0$ emerges automatically within the context of a more general theory.

Acknowledgements

This work was supported by the Robert A. Welch Foundation.

References

1. R.E. Allen, *Int. J. Mod. Phys. A* **12**, 2385 (1997), hep-th/9612041.
2. R.E. Allen, astro-ph/9803079.
3. The cosmological instanton introduced in Refs. 1 and 2 is, of course, unrelated to the one subsequently proposed by S.W. Hawking and N. Turok, hep-th/9802030.
4. A.L. Fetter and J.D. Walecka, *Quantum Theory of Many-Particle Systems* (McGraw-Hill, San Francisco, 1971).
5. M.S. Turner, *Nucl. Phys. A* **621**, 522c (1997), astro-ph/9704024.
6. See, e.g., E.W. Kolb, *Astrophys. J.* **344**, 543 (1989).
7. See, e.g., M. Özer and M.O. Taha, *Nuclear Physics B* **287**, 776 (1987).